

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 209/42, A61K 31/40, C07D 409/12, 471/04, 401/12		A1	(11) International Publication Number: WO 97/34870
			(43) International Publication Date: 25 September 1997 (25.09.97)
(21) International Application Number: PCT/US97/04623		(74) Agent: SARUSSI, Steven, J.; McDonnell Boehnen Hulbert & Berghoff, 300 South Wacker Drive, Chicago, IL 60606 (US).	
(22) International Filing Date: 20 March 1997 (20.03.97)			
(30) Priority Data: 08/620,939 22 March 1996 (22.03.96) US		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(60) Parent Application or Grant (63) Related by Continuation US Filed on 08/620,939 (CIP) 22 March 1996 (22.03.96)		Published With international search report.	
(71) Applicant (for all designated States except US): NEUROGEN CORPORATION [US/US]; 35 Northeast Industrial Road, Branford, CT 06405 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): ALBAUGH, Pamela [US/US]; 81 Long Hill Road, Clinton, CT 06413 (US). LIU, Gang [CN/US]; 144 Montoya Circle, Branford, CT 06405 (US). HUTCHISON, Alan [US/US]; 175 Bartlett Drive, Madison, CT 06443 (US).			

(54) Title: CERTAIN FUSED PYRROLECARBOXAMIDES AS GABA BRAIN RECEPTOR LIGANDS

(57) Abstract

Disclosed are compounds of formula (I), or the pharmaceutically acceptable non-toxic salts thereof, wherein W represents substituted or unsubstituted phenyl; T is hydrogen, halogen, hydroxyl, amino or alkyl; X is hydrogen, hydroxy, or lower alkyl; m is 0, 1, or 2; n is 0, 1, or 2; and R₃ and R₄ represent substituted or unsubstituted organic residues. These compounds are highly selective agonists, antagonists or inverse agonists for GABA_A brain receptors or prodrugs of agonists, antagonists or inverse agonists for GABA_A brain receptors. These compounds are useful in the diagnosis and treatment of anxiety, sleep and seizure disorders, overdose with benzodiazepine drugs and for enhancement of memory.



12/14
us 65
6353109
same disclosure
as 98/02420

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

CERTAIN FUSED PYRROLECARBOXAMIDES AS GABA BRAIN RECEPTOR LIGANDS.

BACKGROUND OF THE INVENTION

5

Field of the Invention

This invention relates to certain fused pyrrolecarboxanilides which selectively bind to GABA_A receptors. This invention also relates to pharmaceutical compositions comprising such compounds. It further relates to the use of such compounds in treating anxiety, sleep and seizure disorders, and overdoses of benzodiazepine-type drugs, and enhancing alertness.

10

Description of the Related Art

γ -Aminobutyric acid (GABA) is regarded as one of the major inhibitory amino acid transmitters in the mammalian brain. Over 30 years have elapsed since its presence in the brain was demonstrated (Roberts & Frankel, J. Biol. Chem. 187: 55-63, 1950; Udenfriend, J. Biol. Chem. 187: 65-69, 1950). Since that time, an enormous amount of effort has been devoted to implicating GABA in the etiology of seizure disorders, sleep, anxiety and cognition (Tallman and Gallager, Ann. Rev. Neuroscience 8: 21-44, 1985). Widely, although unequally, distributed through the mammalian brain, GABA is said to be a transmitter at approximately 30% of the synapses in the brain. In most regions of the brain, GABA is associated with local inhibitory neurons and only in two regions is GABA associated with longer projections. GABA mediates many of its actions through a complex of proteins localized both on cell bodies and nerve endings; these are called GABA_A receptors. Postsynaptic responses to GABA are mediated through alterations in chloride conductance that generally, although not invariably, lead to hyperpolarization of the cell. Recent investigations have indicated that the complex of proteins associated with postsynaptic GABA responses is a major site of action for a number of structurally unrelated compounds capable of modifying postsynaptic responses to GABA. Depending on the mode of interaction, these compounds are capable of producing a spectrum of activities (either sedative, anxiolytic, and anticonvulsant, or wakefulness, seizures, and anxiety).

15

20

25

1,4-Benzodiazepines continue to be among the most widely used drugs in the world. Principal among the benzodiazepines marketed are chlordiazepoxide, diazepam, flurazepam, and triazolam. These compounds are widely used as anxiolytics, sedative-hypnotics, muscle relaxants, and anticonvulsants. A number of these compounds are extremely potent drugs; such potency indicates a site of action with a high affinity and specificity for individual receptors. Early electrophysiological studies indicated that a major action of benzodiazepines was enhancement of GABAergic inhibition. The benzodiazepines were capable of enhancing presynaptic inhibition of a monosynaptic ventral root reflex, a GABA-mediated event (Schmidt et al., 1967, Arch. Exp. Path. Pharmacol. 258: 69-82). All subsequent electrophysiological studies (reviewed in Tallman et al. 1980, Science 207: 274-81, Haefley et al., 1981, Handb. Exptl. Pharmacol. 33: 95-102) have generally confirmed this finding, and by the mid-1970s, there was a general consensus among electrophysiologists that the benzodiazepines could enhance the actions of GABA.

With the discovery of the "receptor" for the benzodiazepines and the subsequent definition of the nature of the interaction between GABA and the benzodiazepines, it appears that the behaviorally important interactions of the benzodiazepines with different neurotransmitter systems are due in a large part to the enhanced ability of GABA itself to modify these systems. Each modified system, in turn, may be associated with the expression of a behavior.

Studies on the mechanistic nature of these interactions depended on the demonstration of a high-affinity benzodiazepine binding site (receptor). Such a receptor is present in the CNS of all vertebrates phylogenetically newer than the boney fishes (Squires & Braestrup 1977, Nature 166: 732-34, Mohler & Okada, 1977, Science 198: 854-51, Mohler & Okada, 1977, Br. J. Psychiatry 133: 261-68). By using tritiated diazepam, and a variety of other compounds, it has been demonstrated that these benzodiazepine binding sites fulfill many of the criteria of pharmacological receptors; binding to these sites *in vitro* is rapid, reversible, stereospecific, and saturable. More importantly, highly significant correlations have been shown between the ability of benzodiazepines to displace diazepam from its binding site and activity in a number of animal behavioral tests predictive of benzodiazepine potency

(Braestrup & Squires 1978, Br. J. Psychiatry 133: 249-60, Mohler & Okada. 1977, Science 198: 854-51, Mohler & Okada. 1977, Br. J. Psychiatry 133: 261-68). The average therapeutic doses of these drugs in man also correlate with receptor potency (Tallman et al. 1980, Science 207: 274-281).

5 In 1978, it became clear that GABA and related analogs could interact at the low affinity (1 mM) GABA binding site to enhance the binding of benzodiazepines to the clonazepam-sensitive site (Tallman et al. 1978, Nature, 274: 383-85). This enhancement was caused by an increase in the affinity of the benzodiazepine binding site due to occupancy of the GABA site. The data were interpreted to mean that both GABA and benzodiazepine sites
10 were allosterically linked in the membrane as part of a complex of proteins. For a number of GABA analogs, the ability to enhance diazepam binding by 50% of maximum and the ability to inhibit the binding of GABA to brain membranes by 50% could be directly correlated. Enhancement of benzodiazepine binding by GABA agonists is blocked by the GABA receptor antagonist (+) bicuculline; the stereoisomer (-) bicuculline is much less active (Tallman et al.,
15 1978, Nature, 274: 383-85).

Soon after the discovery of high affinity binding sites for the benzodiazepines, it was discovered that a triazolopyridazine could interact with benzodiazepine receptors in a number of regions of the brain in a manner consistent with receptor heterogeneity or negative cooperativity. In these studies, Hill coefficients significantly less than one were observed in a
20 number of brain regions, including cortex, hippocampus, and striatum. In cerebellum, triazolopyridazine interacted with benzodiazepine sites with a Hill coefficient of 1 (Squires et al., 1979, Pharma. Biochem. Behav. 10: 825-30, Klepner et al. 1979, Pharmacol. Biochem. Behav. 11: 457-62). Thus, multiple benzodiazepine receptors were predicted in the cortex, hippocampus, striatum, but not in the cerebellum.

25 Based on these studies, extensive receptor autoradiographic localization studies were carried out at a light microscopic level. Although receptor heterogeneity has been demonstrated (Young & Kuhar 1980, J. Pharmacol. Exp. Ther. 212: 337-46, Young et al., 1981 J. Pharmacol Exp. ther 216: 425-430, Niehoff et al. 1982, J. Pharmacol. Exp. Ther. 221: 670-75), no simple correlation between localization of receptor subtypes and the behaviors

associated with the region has emerged from the early studies. In addition, in the cerebellum, where one receptor was predicted from binding studies, autoradiography revealed heterogeneity of receptors (Niedhoff et al., 1982, *J. Pharmacol. Exp. Ther.* 221: 670-75).

A physical basis for the differences in drug specificity for the two apparent subtypes of benzodiazepine sites has been demonstrated by Sieghart & Karobath, 1980, *Nature* 286: 285-87. Using gel electrophoresis in the presence of sodium dodecyl sulfate, the presence of several molecular weight receptors for the benzodiazepines has been reported. The receptors were identified by the covalent incorporation of radioactive flunitrazepam, a benzodiazepine which can covalently label all receptor types. The major labeled bands have molecular weights of 50,000 to 53,000, 55,000, and 57,000 and the triazolopyridazines inhibit labeling of the slightly higher molecular weight forms (53,000, 55,000, 57,000) (Sieghart et al. 1983, *Eur. J. Pharmacol.* 88: 291-99).

At that time, the possibility was raised that the multiple forms of the receptor represent "isoreceptors" or multiple allelic forms of the receptor (Tallman & Gallager 1985, *Ann. Rev. Neurosci.* 8, 21-44). Although common for enzymes, genetically distinct forms of receptors have not generally been described. As we begin to study receptors using specific radioactive probes and electrophoretic techniques, it is almost certain that isoreceptors will emerge as important in investigations of the etiology of psychiatric disorders in people.

The GABA_A receptor subunits have been cloned from bovine and human cDNA libraries (Schoenfield et al., 1988; Duman et al., 1989). A number of distinct cDNAs were identified as subunits of the GABA_A receptor complex by cloning and expression. These are categorized into α , β , γ , δ , ϵ , and provide a molecular basis for the GABA_A receptor heterogeneity and distinctive regional pharmacology (Shivvers et al., 1980; Levitan et al., 1989). The γ subunit appears to enable drugs like benzodiazepines to modify the GABA responses (Pritchett et al., 1989). The presence of low Hill coefficients in the binding of ligands to the GABA_A receptor indicates unique profiles of subtype specific pharmacological action.

Drugs that interact at the GABA_A receptor can possess a spectrum of pharmacological activities depending on their abilities to modify the actions of GABA. For example, the beta-

carbolines were first isolated based upon their ability to inhibit competitively the binding of diazepam to its binding site (Nielsen et al., 1979, Life Sci. 25: 679-86). The receptor binding assay is not totally predictive about the biological activity of such compounds; agonists, partial agonists, inverse agonists, and antagonists can inhibit binding. When the beta-carboline structure was determined, it was possible to synthesize a number of analogs and test these compounds behaviorally. It was immediately realized that the beta-carbolines could antagonize the actions of diazepam behaviorally (Tenen & Hirsch, 1980, Nature 288: 609-10). In addition to this antagonism, beta-carbolines possess intrinsic activity of their own opposite to that of the benzodiazepines; they become known as inverse agonists.

10 In addition, a number of other specific antagonists of the benzodiazepine receptor were developed based on their ability to inhibit the binding of benzodiazepines. The best studied of these compounds is an imidazodiazepine (Hunkeler et al., 1981, Nature 290: 514-516). This compound is a high affinity competitive inhibitor of benzodiazepine and beta-carboline binding and is capable of blocking the pharmacological actions of both these classes of
15 compounds. By itself, it possesses little intrinsic pharmacological activity in animals and humans (Hunkeler et al., 1981, Nature 290: 514-16; Darragh et al., 1983, Eur. J. Clin. Pharmacol. 14: 569-70). When a radiolabeled form of this compound was studied (Mohler & Richards, 1981, Nature 294: 763-65), it was demonstrated that this compound would interact with the same number of sites as the benzodiazepines and beta-carbolines, and that the
20 interactions of these compounds were purely competitive. This compound is the ligand of choice for binding to GABA_A receptors because it does not possess receptor subtype specificity and measures each state of the receptor.


The study of the interactions of a wide variety of compounds similar to the above has led to the categorizing of these compounds. Presently, those compounds possessing activity
25 similar to the benzodiazepines are called agonists. Compounds possessing activity opposite to benzodiazepines are called inverse agonists, and the compounds blocking both types of activity have been termed antagonists. This categorization has been developed to emphasize the fact that a wide variety of compounds can produce a spectrum of pharmacological effects, to indicate that compounds can interact at the same receptor to produce opposite effects, and

to indicate that beta-carbolines and antagonists with intrinsic anxiogenic effects are not synonymous.


A biochemical test for the pharmacological and behavioral properties of compounds that interact with the benzodiazepine receptor continues to emphasize the interaction with the GABAergic system. In contrast to the benzodiazepines, which show an increase in their affinity due to GABA (Tallman et al., 1978, *Nature* 274: 383-85, Tallman et al., 1980, *Science* 207: 274-81), compounds with antagonist properties show little GABA shift (i.e., change in receptor affinity due to GABA) (Mohler & Richards 1981, *Nature* 294: 763-65), and the inverse agonists actually show a decrease in affinity due to GABA (Braestrup & Nielson 1981, *Nature* 294: 472-474). Thus, the GABA shift predicts generally the expected behavioral properties of the compounds.

Various compounds have been prepared as benzodiazepine agonists and antagonists. For Example, U.S. Patents Nos. 3,455,943, 4,435,403, 4,596,808, 4,623,649, and 4,719,210, German Patent No. DE 3,246,932, and Liebigs Ann. Chem. 1986, 1749 teach assorted benzodiazepine agonists and antagonists and related anti-depressant and central nervous system active compounds.

U.S. Patent No. 3,455,943 discloses compounds of the formula:




wherein R₁ is a member of the group consisting of hydrogen and lower alkoxy; R₂ is a member of the group consisting of hydrogen and lower alkoxy; R₃ is a member of the group consisting of hydrogen and lower alkyl; and X is a divalent radical selected from the group consisting of



and the non-toxic acid addition salts thereof.


Other references, such as U.S. Patent No. 4,435,403 and German patent DE 3,246,932 disclose compounds containing the following structural skeleton:



5


where A is carbon or nitrogen.

A variety of indole-3-carboxamides are described in the literature. For example, J. Org. Chem., 42: 1883-1885 (1977) discloses the following compounds.




10

J. Heterocyclic Chem., 14: 519-520 (1977) discloses a compound of the following formula:



None of these indole-3-carboxamides includes an oxy substituent at the 4-position of the indole ring.

5 U.S. Patent No. 5,484,944, the disclosure of which is incorporated herein in its entirety, discloses compounds of the general formula:




or the pharmaceutically acceptable non-toxic salts thereof wherein:


10 T is halogen, hydrogen, hydroxyl, amino or straight or branched chain lower alkoxy having 1-6 carbon atoms;

X is hydrogen, hydroxyl or straight or branched chain lower alkyl having 1-6 carbon atoms;

15 W is phenyl, 2- or 3-thienyl, 2-, 3-, or 4-pyridyl or 6-quinolinyl, each of which may be mono or disubstituted with halogen, cyano, hydroxy, straight or branched chain lower alkyl having 1-6 carbon atoms, amino, mono or dialkylamino where each alkyl is independently straight or branched chain lower alkyl having 1-6 carbon atoms, straight or branched chain lower alkoxy having 1-6 carbon atoms, or NR_1COR_2 , COR_2 , CONR_1R_2 or CO_2R_2 where R_1 and R_2 are the same or different and represent hydrogen or straight or branched chain lower
20 alkyl having 1-6 carbon atoms; and



represents



wherein:

5

Y represents nitrogen or C-R₄;

Z represents N-R₇ or a carbon atom substituted with R₈ and R₉, i.e., C(R₈)(R₉);

n is 1, 2, 3, or 4;

10

R₃ is hydrogen, phenyl, 2-, 3-, or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2-, 3-, or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms;

R₄ is halogen or trifluoromethyl; or

15

-OR₁₀, -COR₁₀, -CO₂R₁₀, -OCOR₁₀, or -R₁₀, where R₁₀ is hydrogen, phenyl, 2-, 3-, or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2-, 3-, or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; or

20

-CONR₁₁R₁₂ or -(CH₂)_mNR₁₁R₁₂, where m is 0, 1, or 2; R₁₁ represents hydrogen, straight or branched chain lower alkyl having 1-6 carbon atoms; and R₁₂ is hydrogen, phenyl, 2-, 3-, or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2-, 3-, or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; or NR₁₁R₁₂ forms a heterocyclic group which is morpholinyl, piperidinyl, pyrrolidinyl, or N-alkyl piperazinyl;

R₅ and R₆ are the same or different and represent

25

hydrogen, halogen, straight or branched chain lower alkyl having 1-6 carbon atoms, or straight or branched chain lower alkoxy having 1-6 carbon atoms;

R₇ is hydrogen, phenyl, 2-, 3-, or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2-, 3-, or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms;


R₈ is hydrogen or straight or branched chain lower alkyl having 1-6 carbon atoms; and

R₉ is -COR₁₃, -CO₂R₁₃ or -R₁₃, where R₁₃ is hydrogen, phenyl, 2-, 3-, or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2-, 3-, or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; or

-CONR₁₄R₁₅ or -(CH₂)_kNR₁₄R₁₅, where k is 0, 1, or 2; R₁₄

represents hydrogen, straight or branched chain lower alkyl having 1-6 carbon atoms; and R₁₅ is hydrogen, phenyl, 2-, 3-, or 4-pyridyl, straight or branched chain lower alkyl having 1-6 carbon atoms, or phenylalkyl or 2-, 3-, or 4-pyridylalkyl where each alkyl is straight or branched chain lower alkyl having 1-6 carbon atoms; or NR₁₄R₁₅ forms a heterocyclic group which is morpholinyl, piperidinyl, pyrrolidinyl, or N-alkyl piperazinyl.

International Application No. PCT/US94/12300, filed October 26, 1994 and published May 4, 1995, the disclosure of which is incorporated herein in its entirety, also discloses pyrrole derivatives of the general formula described in U.S. Patent No. 5,484,944, i.e.,




The substituents on this general formula are as defined in U.S. Patent No. 5,484,944. In addition, U.S. Application S.N. 08/473,509, filed June 7, 1995, the disclosure of which is incorporated herein in its entirety, discloses compounds of the general formula set forth in U.S. Patent No. 5,484,944.

SUMMARY OF THE INVENTION

This invention provides novel compounds of Formula I which interact with a GABA_A binding site, the benzodiazepine receptor.

The invention provides pharmaceutical compositions comprising compounds of Formula I. The invention also provides compounds useful in the diagnosis and treatment of anxiety, sleep and seizure disorders, overdose with benzodiazepine drugs and for enhancement of memory. Accordingly, a broad embodiment of the invention is directed to compounds of general Formula I:



I

or the pharmaceutically acceptable non-toxic salts thereof wherein:

W is aryl or heteroaryl; and

T is halogen, hydrogen, hydroxyl, amino or straight or branched chain lower alkoxy having 1-6 carbon atoms;

X is hydrogen, hydroxyl or straight or branched chain lower alkyl having 1-6 carbon atoms;

m is 0, 1, or 2;

n is 0, 1, or 2; and

R₃ and R₄ are the same or different and are selected from hydrogen, straight or branched lower alkyl having 1-6 carbon atoms, COR₅ or CO₂R₅ where R₅ is straight or branched lower alkyl having 1-6 carbon atoms or cycloalkyl having 3-7 carbon atoms, CONR₆R₇ where R₆ and R₇ are selected independently from hydrogen, straight or branched chain lower alkyl having 1-6 carbon atoms, cycloalkyl having 3-7 carbon atoms, phenyl, 2-, 3-, or 4-pyridyl, or NR₆R₇ forms a heterocyclic group which is morpholinyl, piperidinyl, pyrrolidinyl, or N-alkyl piperazinyl; or

R₃-R₄ together represent a cyclic moiety having 3-7 carbon atoms; and

where each alkyl substituent in Formula I is optionally substituted with at least one group independently selected from hydroxy, mono- or dialkyl amino, phenyl or pyridyl.

These compounds are highly selective agonists, antagonists or inverse agonists for GABA_A brain receptors or prodrugs of agonists, antagonists or inverse agonists for GABA_A brain receptors. In other words, while the compounds of the invention all interact with GABA_A brain receptors, they do not display identical physiologic activity. Thus, these compounds are useful in the diagnosis and treatment of anxiety, sleep and seizure disorders, overdose with benzodiazepine drugs and for enhancement of memory. For example, these compounds can be used to treat overdoses of benzodiazepine-type drugs as they would competitively bind to the benzodiazepine receptor.

DETAILED DESCRIPTION OF THE INVENTION

As used herein, the term "aryl" refers to aromatic carbocyclic groups having a single ring (e.g., phenyl), multiple rings (e.g., biphenyl), or multiple condensed rings in which at least one is aromatic. (e.g., 1,2,3,4-tetrahydronaphthyl, naphthyl, anthryl, or phenanthryl),
 5 which can optionally be substituted with e.g., halogen, lower alkyl, lower alkylthio, trifluoromethyl, lower acyloxy, aryl, and heteroaryl.

A preferred aryl group is phenyl optionally substituted with up to five groups selected independently from halogen, cyano, hydroxy, straight or branched chain lower alkyl having 1-6 carbon atoms or cycloalkyl having 3-7 carbon atoms, amino, mono or dialkylamino where
 10 each alkyl is independently straight or branched chain lower alkyl having 1-6 carbon atoms or cycloalkyl having 3-7 carbon atoms, straight or branched chain lower alkoxy having 1-6 carbon atoms, cycloalkyl alkoxy having 3-7 carbon atoms, or NR_1COR_2 , COR_2 , CONR_1R_2 or CO_2R_2 where R_1 and R_2 are the same or different and represent hydrogen or straight or branched chain lower alkyl having 1-6 carbon atoms or cycloalkyl having 3-7 carbon atoms


15 By heteroaryl is meant aromatic ring systems having at least one and up to four hetero atoms selected from the group consisting of nitrogen, oxygen and sulfur. Examples of heteroaryl groups are pyridyl, pyrimidinyl, pyrrolyl, pyrazolyl, pyrazinyl, pyridazinyl, oxazolyl, naphthyridinyl, isoxazolyl, phthalazinyl, furanyl, quinolinyl, isoquinolinyl, thiazolyl, and thienyl, which can optionally be substituted with, e.g., halogen, lower alkyl, lower alkoxy,
 20 lower alkylthio, trifluoromethyl, lower acyloxy, aryl, heteroaryl, and hydroxy.

The aryl and heteroaryl groups herein are systems characterized by $4n+2$ π electrons, where n is an integer..

In addition to those mentioned above, other examples of the aryl and heteroaryl groups encompassed within the invention are the following:



25



5 As noted above, each of these groups can optionally be mono- or polysubstituted with groups selected independently from, for example, halogen, lower alkyl, lower alkoxy, lower alkylthio, trifluoromethyl, lower acyloxy, aryl, heteroaryl, and hydroxy.


Still other examples of various aryl and heteroaryl groups are shown in Chart D of published International Application WO 93/17025.

10 By "alkyl" and "lower alkyl" in the present invention is meant straight or branched chain alkyl groups having 1-6 carbon atoms, such as, for example, methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, 2-pentyl, isopentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl, and 3-methylpentyl. Unless indicated otherwise, the alkyl group substituents herein are optionally substituted with at least one group independently selected
15 from hydroxy, mono- or dialkyl amino, phenyl or pyridyl.

Where R₁ and R₂ are both alkyl, each alkyl is independently selected from lower alkyl.

By "alkoxy" and "lower alkoxy" in the present invention is meant straight or branched chain alkoxy groups having 1-6 carbon atoms, such as, for example, methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, sec-butoxy, tert-butoxy, pentoxy, 2-pentyl, isopentoxy, neopentoxy, hexoxy, 2-hexoxy, 3-hexoxy, and 3-methylpentoxy.
20


As used herein "cycloalkyl alkoxy" refers to groups of the formula



where a is an integer of from 2 to 6; R' and R'' independently represent hydrogen or alkyl; and b is an integer of from 1 to 6.

By the term "halogen" in the present invention is meant fluorine, bromine, chlorine, and iodine.

By "N-alkylpiperazyl" in the invention is meant radicals of the formula:




where R is alkyl as defined above.

By "monoalkylamino" as used herein is meant an amino substituent substituted with one (1) alkyl group where the alkyl group is lower alkyl as defined above or cycloalkyl having from 3-7 carbon atoms.

By "dialkylamino" as used herein is meant an amino substituent substituted with two (2) alkyl groups where the alkyl groups are independently lower alkyl groups as defined above or cycloalkyl groups having from 3-7 carbon atoms.

The novel compounds encompassed by the instant invention can be described by general formula I set forth above or the pharmaceutically acceptable non-toxic salts thereof.

In addition, the present invention encompasses compounds of Formula II.



wherein

R₃ and R₄ are the same or different and represent hydrogen, alkyl, COR₅ or CO₂R₅ where

R₅ is alkyl or cycloalkyl having 3-7 carbon atoms. CONR₆R₇ where R₆ and R₇ are selected independently from hydrogen, alkyl, cycloalkyl having 3-7 carbon atoms, phenyl, 2-3- or 4-pyridyl, or NR₆R₇ forms a heterocyclic group which is

morpholinyl, piperidinyl, pyrrolidinyl, or N-alkyl piperazinyl; or

R₃-R₄ together represent a cyclic moiety having 3-7 carbon atoms;

R₈ is hydrogen, halogen, hydroxyl, alkyl, alkoxy, cycloalkyl alkoxy having 3-7 carbon atoms, amino, mono- or dialkylamino; and

R₉ is hydrogen, halogen, cyano, hydroxy, alkyl, alkoxy, cycloalkyl alkoxy having 3-7 carbon atoms, amino, mono- or dialkylamino, NR₁COR₂, COR₂, or CO₂R₂ where R₁ and R₂ are the same or different and represent hydrogen, alkyl, or cycloalkyl having 3-7 carbon atoms; and


R₁₀ is hydrogen, halogen, hydroxyl, alkyl, alkoxy, amino, mono- or dialkylamino;

m is 0, 1, or 2; and

n is 0, 1, or 2.

Preferred compounds of Formula II are those where the phenyl group is mono-, di-, or trisubstituted in the 2,4, and/or 5 positions relative to the point of attachment of the phenyl ring to the amide nitrogen.

In addition, the present invention encompasses compounds of Formula III.



wherein

R₃ and R₄ are the same or different and represent hydrogen or alkyl;


R₈ is hydrogen, halogen, hydroxyl, alkyl, alkoxy, cycloalkyl alkoxy having 3-7 carbon atoms, amino, mono- or dialkylamino; and

R₉ is hydrogen, halogen, cyano, hydroxy, alkyl, alkoxy, cycloalkyl alkoxy having 3-7 carbon atoms, amino, mono- or dialkylamino, NR₁COR₂, COR₂, or CO₂R₂ where R₁ and R₂ are the same or different and represent hydrogen, alkyl, or cycloalkyl having 3-7 carbon atoms; and

R₁₀ is hydrogen, halogen, hydroxyl, alkyl, alkoxy, amino, mono- or dialkylamino.

Preferred compounds of Formula III are those where the phenyl group is mono-, di-, or trisubstituted in the 2,4, and/or 5 positions relative to the point of attachment of the phenyl ring to the amide nitrogen. Particularly preferred compounds of Formula III are those where the phenyl group is trisubstituted in the 2,4, and 5 positions relative to the point of attachment of the phenyl ring to the amide nitrogen, and R₈, R₉, and R₁₀ are independently selected from hydrogen, halogen, hydroxy, alkoxy, and alkyl, provided that not all of R₈, R₉, and R₁₀ are hydrogen.

In addition, the present invention encompasses compounds of Formula IV.



IV

wherein

R₃ represents alkyl;

R₈ is hydrogen, halogen, hydroxyl, alkyl, alkoxy, cycloalkyl alkoxy having 3-7 carbon atoms, amino, mono- or dialkylamino; and


R₉ is hydrogen, halogen, cyano, hydroxy, alkyl, alkoxy, cycloalkyl alkoxy having 3-7 carbon atoms, amino, mono- or dialkylamino, NR₁COR₂, COR₂, or CO₂R₂ where R₁ and

R₂ are the same or different and represent hydrogen, alkyl, or cycloalkyl having 3-7 carbon atoms; and

R₁₀ is hydrogen, halogen, hydroxyl, alkyl, alkoxy, amino, mono- or dialkylamino.

Preferred compounds of Formula IV are those where R₁, R₆, and R₁₀ are independently selected from hydrogen, halogen, hydroxy, alkoxy, and alkyl, provided that not all of R₁, R₆, and R₁₀ are hydrogen.

In addition, the present invention encompasses compounds of Formula V.



wherein

R₃ and R₄ are the same or different and represent alkyl;


R₈ is hydrogen, halogen, hydroxyl, alkyl, alkoxy, cycloalkyl alkoxy having 3-7 carbon atoms, amino, mono- or dialkylamino; and

R₉ is hydrogen, halogen, cyano, hydroxy, alkyl, alkoxy, cycloalkyl alkoxy having 3-7 carbon atoms, amino, mono- or dialkylamino, NR₁COR₂, COR₂, or CO₂R₂ where R₁ and R₂ are the same or different and represent hydrogen, alkyl, or cycloalkyl having 3-7 carbon atoms; and

R₁₀ is hydrogen, halogen, hydroxyl, alkyl, alkoxy, amino, mono- or dialkylamino.

Preferred compounds of Formula V are those where R₈, R₉, and R₁₀ are independently selected from hydrogen, halogen, hydroxy, alkoxy, and alkyl, provided that not all of R₈, R₉, and R₁₀ are hydrogen. Particularly preferred compounds of Formula V are those where R₃ and R₄ are both methyl, and R₈, R₉, and R₁₀ are independently selected from hydrogen, halogen, hydroxy, alkoxy, and alkyl, provided that not all of R₈, R₉, and R₁₀ are hydrogen.

In addition, the present invention encompasses compounds of Formula VI



wherein

q is an integer of from 2-6;

R₈ is hydrogen, halogen, hydroxyl, alkyl, alkoxy, cycloalkyl alkoxy having 3-7 carbon atoms, amino, mono- or dialkylamino; and

R₉ is hydrogen, halogen, cyano, hydroxy, alkyl, alkoxy, cycloalkyl alkoxy having 3-7 carbon atoms, amino, mono- or dialkylamino, NR₁COR₂, COR₂, or CO₂R₂ where R₁ and R₂ are the same or different and represent hydrogen, alkyl, or cycloalkyl having 3-7 carbon atoms; and


R₁₀ is hydrogen, halogen, hydroxyl, alkyl, alkoxy, amino, mono- or dialkylamino;

m is 0, 1, or 2; and

n is 0, 1, or 2.

Preferred compounds of Formula VI are those where R₈, R₉, and R₁₀ are independently selected from hydrogen, halogen, hydroxy, alkoxy, and alkyl, provided that not all of R₈, R₉, and R₁₀ are hydrogen.

In addition, the present invention encompasses compounds of Formula VII.



VII


where


G represents aryl or heteroaryl such as, for example, thienyl, thiazolyl, pyridyl, naphthyridinyl,
 5 quinolinyl, or phenyl, each of which is optionally mono-, di- or trisubstituted with
 halogen alkyl alkoxy, or hydroxy; and

R3 and R4 are the same or different and represent hydrogen or alkyl, provided that not both
 R3 and R4 are hydrogen.

Preferred compounds of Formula VII are those where R3 and R4 are C₁₋₃ alkyl, and
 10 more preferably methyl. Other preferred compounds of Formula VII are those where R3 is
 hydrogen and R4 is C₁₋₃ alkyl, and more preferably R4 is methyl.

Preferred compounds of Formula VII include a G group selected from the following:





In the above G groups, the following definitions apply:

R_a is halogen;

R_b is hydroxy;

5 R_c represents alkoxy;

R_d represents alkyl;

R_e represents hydrogen or R_d ;

R_f represents hydrogen, or R_c ; and


R_g represents hydrogen, R_a or R_c .

10 In those formulas where more than one of the same substituent appears, those substituents are the same or different.


Particularly preferred R_a groups in G are fluorine. Particularly preferred R_c groups in G are methoxy and ethoxy. Particularly preferred R_d groups in G are methyl and ethyl.

Representative compounds of the invention are shown below in Table 1.


Table 1




Compound 1




Compound 2




Compound 3




Compound 4




Compound 5




Compound 6




Compound 7



Compound 8




Compound 9



Compound 10

The following numbering system is used to identify positions on the pyrrole ring portion of the compounds of the invention:



Representative compounds of the present invention, which are encompassed by Formula I, include, but are not limited to the compounds in Table I and their pharmaceutically acceptable salts. Non-toxic pharmaceutically acceptable salts include salts of acids such as hydrochloric, phosphoric, hydrobromic, sulfuric, sulfinic, formic, toluenesulfonic, methanesulfonic, nitric, benzoic, citric, tartaric, maleic, hydroiodic, alkanolic such as acetic, $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$ where n is 0-4, and the like. Those skilled in the art will recognize a wide variety of non-toxic pharmaceutically acceptable addition salts.

The present invention also encompasses the prodrugs, preferably acylated prodrugs, of the compounds of Formula I. Those skilled in the art will recognize various synthetic methodologies which may be employed to prepare non-toxic pharmaceutically acceptable addition salts and acylated prodrugs of the compounds encompassed by Formula I.

The pharmaceutical utility of compounds of this invention are indicated by the following assay for GABA_A receptor binding activity.

Assays are carried out as described in Thomas and Tallman (*J. Bio. Chem.* 156: 9838-9842, *J. Neurosci.* 3: 433-440, 1983). Rat cortical tissue is dissected and homogenized in 25 volumes (w/v) of 0.05 M Tris HCl buffer (pH 7.4 at 4°C). The tissue homogenate is centrifuged in the cold (4°C) at 20,000 x g for 20'. The supernatant is decanted and the pellet is rehomogenized in the same volume of buffer and again centrifuged at 20,000 x g. The supernatant is decanted and the pellet is frozen at -20°C overnight. The pellet is then thawed and rehomogenized in 25 volume (original wt/vol) of buffer and the procedure is carried out

twice. The pellet is finally resuspended in 50 volumes (w/vol) of 0.05 M Tris HCl buffer (pH 7.4 at 40°C).

- Incubations contain 100 ml of tissue homogenate, 100 ml of radioligand 0.5 nM (³H-RO15-1788 [³H-Flumazenil] specific activity 80 Ci/mmol), drug or blocker and buffer to a total volume of 500 ml. Incubations are carried for 30 min at 40°C then are rapidly filtered through GFB filters to separate free and bound ligand. Filters are washed twice with fresh 0.05 M Tris HCl buffer (pH 7.4 at 40°C) and counted in a liquid scintillation counter. 1.0 mM diazepam is added to some tubes to determine nonspecific binding. Data are collected in triplicate determinations, averaged and % inhibition of total specific binding is calculated.
- Total Specific Binding = Total - Nonspecific. In some cases, the amounts of unlabeled drugs is varied and total displacement curves of binding are carried out. Data are converted to K_i's; results for compounds of this invention are listed in Table 2.

TABLE 2

Compound Number	K _i (nM)
1	21
2	12
3	13
4	1
5	0.5
6	1
7	10
8	80
9	4
10	12

- The compounds of general formula I may be administered orally, topically, parenterally, by inhalation or spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition, there is provided a pharmaceutical

formulation comprising a compound of general formula I and a pharmaceutically acceptable carrier. One or more compounds of general formula I may be present in association with one or more non-toxic pharmaceutically acceptable carriers and/or diluents and/or adjuvants and if desired other active ingredients. The pharmaceutical compositions containing compounds of
5 general formula I may be in a form suitable for oral use. for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs.

Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may
10 contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium
15 carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a
20 longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with
25 water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting

agents may be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters
5 derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more
10 sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and
15 flavoring agents may be added to provide palatable oral preparations. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents
20 and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

Pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may
25 be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be
5 formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride
10 solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono-or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compounds of general formula I may also be administered in the form of
15 suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.


Compounds of general formula I may be administered parenterally in a sterile
20 medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anaesthetics, preservatives and buffering agents can be dissolved in the vehicle.

Dosage levels of the order of from about 0.1 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 0.5 mg to
25 about 7 g per patient per day). The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Dosage unit forms will generally contain between from about 1 mg to about 500 mg of an active ingredient.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular
5 disease undergoing therapy.

An illustration of the preparation of compounds of the present invention is given in Scheme I.

Scheme I



where W, m, n, R₃, and R₄ are defined as above.

- 5 Those having skill in the art will recognize that the starting materials may be varied and additional steps employed to produce compounds encompassed by the present invention, as demonstrated by the following examples. In some cases protection of certain reactive functionalities may be necessary to achieve some of the above transformations. In general the

need for such protecting groups will be apparent to those skilled in the art of organic synthesis as well as the conditions necessary to attach and remove such groups.

The disclosures in this application of all articles and references, including patents, are incorporated herein by reference in their entirety.

5 The invention is illustrated further by the following examples which are not to be construed as limiting the invention in scope or spirit to the specific procedures described in them.


Example 1

Preparation of starting materials and intermediates

The starting materials and various intermediates may be obtained from commercial sources, prepared from commercially available organic compounds, or prepared using well known synthetic methods.

Representative examples of methods for preparing intermediates of the invention are set forth below.

1. Ethyl 3-hydroxy 4-oxo-6-methyl-2,3,4,5,6,7-hexahydrobenzofuran-3-carboxylate




10

To a stirred mixture of 5-methyl-1,3-cyclohexanedione (10.25 g, 81 mmol) and potassium carbonate (22.46 g, 162 mmol) in dichloromethane (200 mL) at 0°C was added a solution of ethyl bromopyruvate (10.7 mL, 85 mmol) in dichloromethane (50 mL). The reaction was allowed to reach ambient temperature, stirred for 18 hours, then poured into saturated aqueous ammonium chloride. After adjusting to neutral pH with aqueous hydrochloric acid, the mixture was extracted 2X with dichloromethane, the combined organic layers were dried over sodium sulfate, filtered, and concentrated *in vacuo* to give ethyl 3-hydroxy 4-oxo-6-methyl-2,3,4,5,6,7-hexahydrobenzofuran-3-carboxylate (18.48 g).

20

2. Ethyl 4-oxo-6-methyl-4,5,6,7-tetrahydrobenzofuran-3-carboxylate




A solution of methanesulfonyl chloride (6.1 mL, 78.5 mmol) in dichloromethane (50 mL) was added to a stirring solution of ethyl 3-hydroxy 4-oxo-6-methyl-2,3,4,5,6,7-hexahydrobenzofuran-3-carboxylic acid (18.48 g, 76 mmol) and triethylamine (21.4 mL, 154 mmol) in dichloromethane (150 mL) at 0°C. The mixture was allowed to reach ambient

25

temperature, stirred for 2 hours, then poured into aqueous 1N sodium hydroxide and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and concentrated in vacuo to give ethyl 4-oxo-6-methyl-4,5,6,7-tetrahydrobenzofuran-3-carboxylate (16.86 g).

5


3. 4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxylic acid



A stirred mixture of ethyl 4-oxo-6-methyl-4,5,6,7-tetrahydrobenzofuran-3-carboxylate (15.7 g, 71 mmol) and ammonium acetate (9.54 g, 124 mmol) in N,N-dimethylformamide (75 mL) was heated at 100°C for 2 hours. The reaction mixture was concentrated *in vacuo*, ice water was added, and the precipitate collected, rinsed with water then diethyl ether and dried to give ethyl 4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxylate (5.94 g). To this ester was added aqueous 5N sodium hydroxide (50 mL) and ethyl alcohol (10 mL) and the mixture heated at reflux for 40 minutes. The reaction mixture was cooled in an ice water bath, acidified with aqueous hydrochloric acid, and the precipitate collected, rinsed with water then diethyl ether and dried to give 4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxylic acid (5.2 g). m.p. 210-211°C.

20 4. 4-Oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxylic acid (m.p. 231-232°C) was prepared essentially according to the procedures described in Parts 1-3 of this examples.

Example 2



To a stirred solution of 4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxylic acid (155 mg, 0.75 mmol) and triethylamine (209 μ L, 1.5 mmol) in dimethylformamide (4 mL) at 0°C was added ethyl chloroformate (143 μ L, 1.5 mmol). After stirring an additional 45 minutes, 2-fluoroaniline (145 μ L, 1.5 mmol) was added. The reaction mixture was stirred for 30 minutes, then poured into aqueous 3.6N hydrochloric acid and extracted 2X with ethyl acetate. The combined organic layers were washed with water, dried over magnesium sulfate, filtered, and concentrated *in vacuo*. To the residue was added aqueous 5N sodium hydroxide (5 mL) and ethyl alcohol (1 mL), and the mixture was heated at reflux for 30 minutes. After cooling in an ice water bath, the reaction mixture was acidified with hydrochloric acid, the precipitate was collected, rinsed with water, and dried to give 55 mg of N-(2-fluorophenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 2).

Example 3

The following compounds are prepared essentially according to the procedures described in Examples 1 and 2.

(a) N-Phenyl-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 1).

(b) N-(2-Fluorophenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 259-261°C.

(c) N-(3-Fluorophenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 268-270°C.

(d) N-(2, 4- Difluorophenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

5 (e) N-(2,4-Difluorophenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide .

(f) N-(3Methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

10

(g) N-(2-Hydroxy-4-methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 190-192°C.

15 (h) N-(3-Hydroxy-4-methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 282-284°C.

(i) N-(2-Fluoro-4-methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 213-215°C.

20 (j) N-(2-Fluoro-4-methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

(k) N-(2-Fluoro-5-methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

25

(l) N-(2-Fluoro-4-methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 225-227°C.

(m) N-(2-Methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 3).

5 (n) N-(4-Ethoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

(o) N-(4-Methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

10 (p) N-(2-Hydroxy-4-methylphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 201-203°C.

(q) N-Phenyl-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 4); mp 278-279°C.

15

(r) N-(2-Fluorophenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 5); mp 264-265°C.

20 (s) N-(3-Fluorophenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 302-303°C.

(t) N-(4-Fluorophenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 262-264°C.

25 (u) N-(3-Methoxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 234-235°C.

(v) N-(4-Hydroxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 320°C.

(w) N-(2-Fluoro-4-hydroxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 330°C.

5 (x) N-(2-Hydroxy-4-methoxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 236-238°C.

(y) N-(4-Methoxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 260-261°C.

10

(z) N-(2-Fluoro-4-methoxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 6); mp 217-219°C.

(aa) N-(4-Ethoxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-
15 carboxamide; mp 269°C.

(bb) N-(2-Fluoro-4-ethoxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 224-225°C.

20 (cc) N-(3,4-Dihydroxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 267-269°C.

(dd) N-(2-Hydroxy-4-methylphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 258-260°C.

25

(ee) N-(3-Thienyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 7).

(ff) N-(2-Thiazoyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.


- (gg) N-(5-Methyl-2-thiazolyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- (hh) N-(3-Pyridyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 237-239°C.
- 5 (ii) N-(4-Methoxy-3-pyridyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 217-218°C.
- (jj) N-(2-Chloro-1,8-naphthyridin-7-yl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 278-280°C.
- (kk) N-(1,8-Napthyridin-2-yl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (compound 8); mp 389-390°C.
- 10 (ll) N-(3-Pyridyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 225-227°C.
- (mm) N-(4-Pyridyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 280-290°C.
- 15 (nn) N-(1,8-Napthyridin-2-yl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 9).
- (oo) N-(6-Methyl-1,8-napthridin-2-yl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; mp 338-340°C(d).
- (pp) N-(2-Quinoliny)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide
- 20 (Compound 10); mp 273-275°C.
- (qq) N-(4-Pyridyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxmide.

The invention and the manner and process of making and using it, are now described in such full, clear, concise and exact terms as to enable any person skilled in the art to which it

25 pertains, to make and use the same. It is to be understood that the foregoing describes preferred embodiments of the present invention and that modifications may be made therein without departing from the spirit or scope of the present invention as set forth in the claims. To particularly point out and distinctly claim the subject matter regarded as invention, the following claims conclude this specification.

WHAT IS CLAIMED IS:


1. A compound of the formula:



or the pharmaceutically acceptable non-toxic salts thereof wherein:

- 5 W is aryl or heteroaryl, each of which is optionally substituted with up to five groups selected independently from
- halogen, cyano, hydroxy, alkyl or cycloalkyl having 3-7 carbon atoms,
- amino or mono- or dialkylamino where each alkyl is independently lower alkyl or
- 10 cycloalkyl having 3-7 carbon atoms,
- alkoxy or cycloalkyl alkoxy having 3-7 carbon atoms, or
- NR₁COR₂, COR₂, CONR₁R₂ or CO₂R₂ where R₁ and R₂ are the same or different
- and represent hydrogen or alkyl or cycloalkyl having 3-7 carbon atoms; and
- T is halogen, hydrogen, hydroxyl, amino or alkoxy;
- X is hydrogen, hydroxyl or alkyl;
- 15 m is 0, 1, or 2;
- n is 0, 1, or 2; and
- R₃ and R₄ are the same or different and represent hydrogen, alkyl, COR₅ or CO₂R₅ where
- R₅ is alkyl or cycloalkyl having 3-7 carbon atoms, CONR₆R₇ where R₆ and R₇ are
- 20 selected independently from hydrogen, alkyl, cycloalkyl having 3-7 carbon atoms,
- phenyl, 2-,3-, or 4-pyridyl, or NR₆R₇ forms a heterocyclic group which is
- morpholinyl, piperidinyl, pyrrolidinyl, or N-alkyl piperazinyl; or
- R₃-R₄ together represent a cyclic moiety having 3-7 carbon atoms.

2. A compound of the formula:



wherein

R3 and R4 are the same or different and represent hydrogen, alkyl, COR5 or CO2R5 where

R5 is alkyl or cycloalkyl having 3-7 carbon atoms, CONR6R7 where R6 and R7 are

5 selected independently from hydrogen, alkyl, cycloalkyl having 3-7 carbon atoms, phenyl, 2-,3-, or 4-pyridyl, or NR6R7 forms a heterocyclic group which is morpholinyl, piperidinyl, pyrrolidinyl, or N-alkyl piperazinyl; or

R3-R4 together represent a cyclic moiety having 3-7 carbon atoms;

R8 is hydrogen, halogen, hydroxyl, alkyl, alkoxy, cycloalkyl alkoxy having 3-7 carbon atoms,

10 amino, mono- or dialkylamino; and

R9 is hydrogen, halogen, cyano, hydroxy, alkyl, alkoxy, cycloalkyl alkoxy having 3-7 carbon atoms, amino, mono- or dialkylamino, NR1COR2, COR2, or CO2R2 where R1 and


R2 are the same or different and represent hydrogen, alkyl, or cycloalkyl having 3-7 carbon atoms; and

15 R10 is hydrogen, halogen, hydroxyl, alkyl, alkoxy, amino, mono- or dialkylamino;

m is 0, 1, or 2; and

n is 0, 1, or 2.

3. A compound of the formula:



20

where

G represents thienyl, thiazolyl, pyridyl, naphthyridinyl, quinolinyl, or phenyl, each of which is optionally mono-, di- or trisubstituted with halogen, alkyl, alkoxy, or hydroxy; and

R₃ and R₄ are the same or different and represent hydrogen or alkyl, provided that not both

5 R₃ and R₄ are hydrogen.

4. A compound according to claim 3, wherein R₃ and R₄ are hydrogen or C₁₋₃ alkyl, provided that not both R₃ and R₄ are hydrogen.

5. A compound according to claim 3, wherein R₃ and R₄ are methyl.

10 6. A compound according to claim 3, wherein R₃ is methyl and R₄ is hydrogen.

7. A compound according to claim 1, which is N-Phenyl-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

8. A compound according to claim 1, which is N-(2-Fluorophenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

15 9. A compound according to claim 1, which is N-(3-Fluorophenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

10. A compound according to claim 1, which is N-(4-Fluorophenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

20 11. A compound according to claim 1, which is N-(2,4-Difluorophenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide .

12. A compound according to claim 1, which is N-(2,6-Difluorophenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide .

13. A compound according to claim 1, which is N-(3-Methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide .

25 14. A compound according to claim 1, which is N-(2-Hydroxy-4-methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

15. A compound according to claim 1, which is N-(3-Hydroxy-4-methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

16. A compound according to claim 1, which is N-(2-Fluoro-4-methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
17. A compound according to claim 1, which is N-(2-Fluoro-4-ethoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 5 18. A compound according to claim 1, which is N-(2-Fluoro-5-methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
19. A compound according to claim 1, which is N-(2-Fluoro-4-hydroxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
20. A compound according to claim 1, which is N-(4-Methoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 10 21. A compound according to claim 1, which is N-(4-Ethoxyphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
22. A compound according to claim 1, which is N-(4-Methylphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 15 23. A compound according to claim 1, which is N-(2-Hydroxy-4-methylphenyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
24. A compound according to claim 1, which is N-Phenyl-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
25. A compound according to claim 1, which is N-(2-Fluorophenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 20 26. A compound according to claim 1, which is N-(3-Fluorophenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
27. A compound according to claim 1, which is N-(4-Fluorophenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 25 28. A compound according to claim 1, which is N-(3-Methoxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
29. A compound according to claim 1, which is N-(4-Hydroxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

30. A compound according to claim 1, which is N-(2-Fluoro-4-hydroxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
31. A compound according to claim 1, which is N-(2-Hydroxy-4-methoxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 5 32. A compound according to claim 1, which is N-(4-Methoxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
33. A compound according to claim 1, which is N-(2-Fluoro-4-methoxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
34. A compound according to claim 1, which is N-(4-Ethoxyphenyl)-4-oxo-6-
10 methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
35. A compound according to claim 1, which is N-(2-Fluoro-4-ethoxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
36. A compound according to claim 1, which is N-(3,4-Dihydroxyphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 15 37. A compound according to claim 1, which is N-(2-Hydroxy-4-methylphenyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
38. A compound according to claim 1, which is N-(3-Thienyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
39. A compound according to claim 1, which is N-(2-Thiazoyl)-4-oxo-6,6-
20 dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
40. A compound according to claim 1, which is N-(5-Methyl-2-thiazolyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
41. A compound according to claim 1, which is N-(3-Pyridyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 25 42. A compound according to claim 1, which is N-(4-Methoxy-3-pyridyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
43. A compound according to claim 1, which is N-(2-Chloro-1,8-naphthyridin-7-yl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

44. A compound according to claim 1, which is N-(1,8-Napthyridin-2-yl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
45. A compound according to claim 1, which is N-(3-Pyridyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
- 5 46. A compound according to claim 1, which is N-(4-Pyridyl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
47. A compound according to claim 1, which is N-(1,8-Napthyridin-2-yl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
48. A compound according to claim 1, which is N-(6-Methyl-1,8-napthridin-2-yl)-
10 4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
49. A compound according to claim 1, which is N-(2-Quinoliny)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.
50. A compound according to claim 1, which is N-(4-Pyridyl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

15

INTERNATIONAL SEARCH REPORT

Int. Application No.
PCT/US 97/04623

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D209/42 A61K31/40 C07D409/12 C07D471/04 C07D401/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 11885 A (NEUROGEN CORPORATION) 4 May 1995 cited in the application * complete document * -----	1-6

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

11 June 1997

Date of mailing of the international search report

18. 06. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC1/US 97/04623

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9511885 A	04-05-95	US 5484944 A	16-01-96
		AU 8126594 A	22-05-95
		CA 2175204 A	11-05-95
		CN 1143953 A	26-02-97
		EP 0725775 A	14-08-96
		FI 961776 A	14-06-96
		NO 961655 A	10-06-96
		PL 314136 A	19-08-96
		SE 9601563 A	22-05-96
		US 5608079 A	04-03-97
